

## BINDING AND CYTOTOXICITY OF CERATO-ULMIN TO ELM CELLS

G. Del Sorbo<sup>1</sup>, P. Ambrosino<sup>1</sup>, C. Comparini<sup>2</sup>, S. Tegli<sup>2</sup>, A. Scala<sup>2</sup> and F. Scala<sup>1</sup>

<sup>1</sup>Department Ar.Bo.Pa.Ve., Section of Plant Pathology,  
University of Naples Federico II, Via Università 100, 80055 Portici (Naples), Italy  
<sup>2</sup>Department of Agricultural Biotechnology, Section of Plant Pathology,  
University of Florence, Piazzale delle Cascine 18, 50144 Florence, Italy

## SUMMARY

*In vitro* cultivated elm cells were used to study the cytotoxicity of cerato-ulmin (= CU), a hydrophobin produced by *Ophiostoma ulmi* and *O. novo-ulmi*, two causal agents of Dutch elm disease (= DED). Purified CU was found to bind to elm cells and cause a dose-dependent death both on cells derived from a DED susceptible (C06) and a moderately resistant (196/6) elm clone. Indirect immunofluorescence (IF) tests, using a rabbit anti-CU antiserum, indicated that homogeneous CU binds to elm but not to cells of fennel and pear, two non-host species of *Ophiostoma*. The binding was demonstrated by quantitative ELISA on extracts from elm cells incubated with purified CU and by measuring the decrease in CU concentration in the incubation mixture. The equation of the saturation binding curve indicated that the maximum quantity of bound CU for elm clones C06 and 196/6 were similar (351.92 and 355.59 pmoles mg<sup>-1</sup> d.wt. elm cells, respectively), as similar were the equilibrium dissociation constants (7.09 and 8.08 µM CU, respectively). The increase in bound CU correlated with the intensity of the fluorescence signal on elm cells and a higher mortality. Elm cell-*Ophiostoma* spp. dual cultures in Takai or Murashige & Skoog-derived media were used to test the ability of elm cells to support growth of *Ophiostoma* fungi and CU production. Cells derived from two elm clones supported both abundant fungal growth and CU production. Fungal growth was supported by cells of the *Ophiostoma* non-hosts fennel and pear, whereas CU was only produced in the presence of fennel cells. The binding and direct cytotoxicity of CU on elm cells support its role as a pathogenicity factor in DED.

*Key words:* *Ophiostoma* spp., *Ulmus* spp., toxin binding, *in vitro* fungus-plant cells dual cultures.

## INTRODUCTION

The Ascomycetes *Ophiostoma ulmi* (Buisman) Nann-

feldt and *Ophiostoma novo-ulmi* Brasier have caused two successive pandemics of a very destructive disease of numerous plant species belonging to genus *Ulmus*, which is known as Dutch elm disease (DED). The second pandemic of DED, caused by *O. novo-ulmi* with the subspecies *novo-ulmi* and *americana* (Brasier and Kirk, 2001), formerly designated Eurasian (=EAN) and North-American (=NAN) races, respectively, was more severe than the first one and almost destroyed the European and North-American native elms.

The phytotoxic hydrophobin cerato-ulmin (CU) (Bowden *et al.*, 1994; Scala *et al.*, 1994; Stringer and Timberlake, 1993; Yaguchi *et al.*, 1993) is a component of the mycelial surface of *O. novo-ulmi*, *O. ulmi* and *Ophiostoma quercus* (Georgevitch), the latter reported as *Ophiostoma piceae* (Munch) H. and P. Sydow (Scala *et al.*, 1997a; Svircev *et al.*, 1988; Takai and Hiratsuka, 1980), and is released in shake liquid culture by only *O. ulmi*, *O. novo-ulmi* and *Ophiostoma himal-ulmi* (Brasier, 1991; Brasier *et al.*, 1990; Brasier and Mehrotra, 1995; Nordin *et al.*, 1987; Richards and Takai, 1973; Scala *et al.*, 1997a). The role of CU in DED pathogenesis is still being debated (Del Sorbo *et al.*, 2002). Many evidences suggest a key role of the toxin in virulence of DED pathogens. For instance, in DED-susceptible elm genotypes, homogeneous CU causes internal and external symptoms identical to those of DED (Takai, 1974; Takai *et al.*, 1983; Takai and Hiratsuka, 1984), alteration of several physiological parameters of both cuttings and leaves (Richards, 1993; Richards and Takai, 1984) and inhibition of division of *in vitro* deep-plated cells (Tegli *et al.*, 1993).

However, there is evidence against a role of CU in virulence. Some naturally-occurring (Brasier *et al.*, 1995) or laboratory-induced (Tegli and Scala, 1996) mutants of *O. novo-ulmi*, which lost the ability to produce CU *in vitro*, maintained their virulence on elm. Transformation-mediated disruption of the CU-encoding gene in *O. novo-ulmi* did not determine changes in virulence on rooted cuttings and seedlings of highly susceptible elm genotypes (Bowden *et al.*, 1996). Transgenic expression of the *cu* gene of *O. novo-ulmi* in *O. ulmi*, a less virulent species, determined an increase in conidial adhesion properties to DED vectors and resistance to desiccation but not an increase in virulence on *O. americana* plants (Temple *et al.*, 1997). Et Touil *et al.*

(1999) reported that virulence of two strains of *O. novo-ulmi*, which produce similar amounts of CU *in vitro*, depends on the single nuclear gene *Pat1*.

CU production by DED pathogens in axenic culture depends on several factors, including fungal genotype. Isolates belonging to both *novo-ulmi* and *americana* subspecies of *O. novo-ulmi* produce more CU than *O. ulmi*, often in amounts significantly correlated to their respective pathogenicity levels on elm (Brasier, 1991; Brasier *et al.*, 1990). Temperature affects CU production, as the two species have two different temperature optima: 21-23°C for *O. novo-ulmi* and 31-33°C for *O. ulmi* (Tegli *et al.*, 1994). CU production is also influenced by the carbon and nitrogen source, as well as by salt concentration of culture medium (Takai, 1978).

The role in pathogenesis of toxins produced *in vitro* by phytopathogenic fungi has widely been discussed. Mitchell (1981) assumed that toxins produced *in vitro* by phytopathogenic fungi are also produced *in planta* during pathogenesis. On the other hand, Gilchrist (1983) suggested that phytotoxic metabolites present in fungal culture filtrates could be culture-derived artifacts, normally not produced during pathogenesis. Yoder (1980; 1981) stated that virulence cannot be always correlated with the quantity of toxin produced in culture. CU was detected in both *O. novo-ulmi* naturally-infected and artificially-inoculated elms (Richards and Takai, 1988). Scala *et al.* (1997b) reported that the amounts of CU produced *in planta* are correlated with those produced *in vitro* by *O. novo-ulmi* and *O. ulmi*. Moreover DED symptoms were associated with high CU contents on leaves. Transgenic expression of the *cu* gene originating from a high CU-producer isolate of *O. novo-ulmi* in *O. quercus*, a saprophyte on elm, enables the fungus to cause typical DED symptoms (Del Sorbo *et al.*, 2000).

In this paper, we studied the effect of the addition of CU to elm cell cultures. This system simulates *in vitro* the condition occurring *in planta* during pathogenesis. The finding that CU binds and determines toxicity to elm cells contributes to a better understanding of the role of CU in DED. We also analysed *in vitro* *Ophiostoma* spp. - elm cell dual cultures with the aim to elucidate whether CU production by the DED fungi is influenced by the presence of host plant cells.

## MATERIALS AND METHODS

**Fungal cultures.** The isolates H328 and H327 of *O. novo-ulmi* ssp. *novo-ulmi*, the isolates 182 and H351 of *O. novo-ulmi* ssp. *americana*, and the isolates E2, R21 and Yu99 of *O. ulmi* are described in previous papers (Scala *et al.*, 1994; Tegli *et al.*, 1994; Tegli and Scala, 1996). The isolate Yu16 of *O. ulmi* was provided by Prof. L. Mitterpergher (Istituto per la Protezione delle Piante, CNR, Firenze, Italy). The isolate H162 of *O. novo-ulmi* ssp. *americana* was from the collection of Prof.

C.M. Brasier (Forest Research Station, Alice Holt Lodge, Farnham, UK). Long- and short-term maintenance of isolates and culture conditions are reported by Scala *et al.* (1994). Budding spores (blastoconidia) were obtained in shake culture in the standard Takai liquid medium (TK) (Takai, 1978) amended according to Scala *et al.* (1994). Blastoconidia concentration was directly estimated by a hemocytometer.

***In vitro* callus cultures and cell suspensions.** *In vitro* elm callus cultures were obtained from two different elm clones, C06 (very susceptible to DED) and 196/6 (moderately resistant to DED). Callus was initiated from young leaves previously surface-sterilised under vacuum in 1% commercial sodium-hypoclorite solution for 30 min and then rinsed four times in sterile distilled water. Leaf segments, including a portion of the mid-vein, were plated on solid Murashige & Skoog medium (MS) (M0404, Sigma-Aldrich, Milan, Italy) supplemented with 2 µM 2,4 dichlorophenoxyacetic acid, 5 µM kinetin, 2 mg ml<sup>-1</sup> glycine, 10 mg ml<sup>-1</sup> ascorbic acid, 30 g l<sup>-1</sup> sucrose and 7 g l<sup>-1</sup> agar. Cultures were incubated in the dark at 23±1°C. Proliferating callus cultures were then routinely transferred to fresh solid MS every 4 weeks. A fine elm cell suspension was obtained by inoculating 10 g of friable callus in 100 ml of liquid MS medium. Agitation of the culture at 100 rpm for 2 days caused the release of single cells and small cell aggregates into the medium. The cell suspension was filtered through a 60 mesh stainless steel screen, washed twice with fresh medium and used for experiments.

Fennel and pear cell suspensions were obtained from actively proliferating callus cultures using the same procedure described for elm cells.

**Cerato-ulmin - elm cell interaction.** Homogeneous CU was obtained from the culture filtrate of the EAN *O. novo-ulmi* H328 (Scala *et al.*, 1994). Interaction between homogeneous CU and elm cells was studied by two different systems. In the first one, different amounts of CU were added to 10 ml aliquots of a standard 5·10<sup>4</sup> ml<sup>-1</sup> cell suspensions of elm clones C06 or 196/6 in order to have a CU concentration range from 0.1 to 32 µM. Elm cell viability and IF assays were performed 3 days after CU administration. Data were collected from two independent experiments, each performed in triplicate. The effect of the same range of CU concentrations was tested in 10 ml suspensions of fennel or pear cells (5·10<sup>4</sup> ml<sup>-1</sup>).

In the second one, cell suspensions of elm clone 196/6 containing increasing cell densities (ranging from 0.25·10<sup>4</sup> to 2·10<sup>4</sup> cells ml<sup>-1</sup>) were incubated in presence of a fixed CU concentration (5 µM) at 23±1°C in agitation (100 rpm). The concentration of CU in solution was measured by indirect ELISA after 1.5, 3 and 6 days of incubation. The experiment was done in triplicate and repeated two times using different batches of cells.

**Evans blue assay.** Elm cell viability was evaluated using the Evans blue stain according to the method of Baker and Mock (1994), with slight modifications. One-ml aliquots of cell suspensions were transferred in test tubes and incubated with 0.5 ml of 0.1% (w/v) aqueous Evans blue for 5 min. Cells were washed in a Sartorius funnel fitted with 3  $\mu\text{m}$  Millipore membranes with distilled water until no further blue eluted from cells. After washing, cells were gently transferred from membranes in 1.5 ml tubes containing 500  $\mu\text{l}$  of 1% (w/v) sodium dodecyl sulphate (SDS) aqueous solution and 100  $\mu\text{g}$  of carborundum. Cells were ground using a micro-pestle, and the homogenate diluted with 500  $\mu\text{l}$  of distilled water. Tubes were then briefly vortexed, incubated for 2 h at room temperature, and centrifuged at 9,000  $g$  for 3 min. A 0.8 ml aliquot of the supernatant was removed, and the absorbance determined spectrophotometrically at 600 nm ( $A_{600}$ ). The freshly prepared cell suspension was assumed to contain 100% of viable cells, and gave a  $A_{600}$  of 0.300. A suspension containing 100% of dead cells, obtained by boiling cells for 10 min, gave a  $A_{600}$  of 0.700. Identical values of  $A_{600}$  were given by an elm cell suspension exposed to lethal concentrations of CU, whose inability to reproduce was checked by plating (Tegli *et al.*, 1993). The percentage of dead cells in each sample was calculated as follows:

$$\% \text{ of dead cells} = \frac{A_{600} \text{ of sample} - 0.300}{0.700 - 0.300} \cdot 100$$

**Immunofluorescence (IF) assay.** Cell suspensions samples (2 ml) were collected and centrifuged for 10 min at 5,000  $g$  at 4°C. The pellet was washed three times with 0.9% (w/v) NaCl and once with distilled water and then resuspended to a concentration of  $10^4$  plant cells  $\text{ml}^{-1}$ . For IF assay, 5  $\mu\text{l}$  of this suspension were dispensed onto 12-well Multitest slides (Flow Laboratories, McLean, VA, USA) and dried at 55°C. Wells were washed with PBS, filled with 20  $\mu\text{l}$  of anti-CU antiserum diluted 1:100 in PBS and incubated for 1 h at room temperature. Control wells were prepared by using preimmune serum diluted 1:100. After three washings with PBS, wells were filled with 20  $\mu\text{l}$  of a 1:40 dilution of goat antirabbit IgG fluorescein isothiocyanate conjugate (Sigma-Aldrich, Milan, Italy) in PBS and incubated for 1 h at room temperature. Slides were washed with distilled water and examined with a Zeiss Axioskop microscope equipped for epifluorescence with an HBO 50 lamp (exciter blue filter, BP 450-490 and barrier filter LP 520).

The intensity of fluorescence was recorded on an arbitrary 0-3 scale, where 0 = no visible fluorescence; 1 = weak fluorescence; 2 = moderate fluorescence; 3 = intense fluorescence.

**Fungus-elm cell suspensions dual cultures.** Fungus-elm cell dual cultures were obtained by inoculating 50

ml Erlenmeyer flasks with 10 ml of a  $5 \cdot 10^4 \text{ ml}^{-1}$  elm cell suspension and  $5 \cdot 10^5 \text{ ml}^{-1}$  blastoconidia. Four different liquid media were used: TK without yeast extract (YE) ( $\text{TK}_0$ ),  $\text{TK}_0$  added with 0.3  $\text{g l}^{-1}$  YE ( $\text{TK}_{0.3\text{YE}}$ ), MS, MS added with 2  $\text{g l}^{-1}$  YE ( $\text{MS}_{2\text{YE}}$ ). Scala (1997) demonstrated that growth of *Ophiostoma* spp. and production of CU in quantities similar to those obtained in the standard medium TK (which contains 2  $\text{g l}^{-1}$  YE), are supported at least by 0.3 and 2  $\text{g l}^{-1}$  YE added to  $\text{TK}_0$  and MS, respectively. Dual cultures were incubated for 10 days in a rotary shaker at 100 rpm at  $23 \pm 1^\circ\text{C}$  in the dark. Dual cultures were also prepared using fennel or pear cells and grown in similar conditions.

**CU assays.** CU concentration was determined by two different assays. CU concentration in liquid media was determined according to the turbidimetric method of Takai and Richards (1978) as modified by Scala *et al.* (1994) and expressed as CU production index (CPI). This index is based on a linear relation between the optical density at 400 nm of a shaken sample and the decimal logarithm of the dilution factor. A CPI value of 100 corresponds to a CU concentration of 26.4  $\mu\text{g ml}^{-1}$  (Scala *et al.*, 1994).

CU bound to elm cells was quantitatively determined by ELISA in cell extracts. An antiserum specific for CU from *O. novo-ulmi* isolate H328 was used (Scala *et al.*, 1994). Elm cell suspensions were collected and centrifuged at 5,000  $g$  for 15 min at 4°C. The pellet was washed three times with 0.9% (w/v) NaCl and once with distilled water, and then freeze-dried. A freeze-dried aliquot (20 mg) of each cell suspension sample was extracted for 2 min with 500  $\mu\text{l}$  of 60% (v/v) ethanol and centrifuged for 5 min at 13,000  $g$ . The supernatant was dried under vacuum. The residue was solubilized in 500  $\mu\text{l}$  phosphate buffered saline 0.15 M, pH 7.2 (PBS) and used for coating of wells of Falcon 3911 Microtest flexible plates (Becton Dickinson Labware, Oxnard, CA, USA). Test samples (50  $\mu\text{l}$ ) were added to triplicate wells for 3 h at 37°C. The wells were washed three times with PBS and any remaining binding sites were blocked with 50  $\mu\text{l}$ /well of 0.5% (w/v) gelatine in PBS for 2 h at 37°C. Each assay included a standard calibration curve that was generated by preparing serial dilutions of homogeneous CU in PBS. Wells coated with 50  $\mu\text{l}$  of extracts from cells inoculated with distilled water served as blanks. Absorbance values for the blanks were subtracted from all test values. Following saturation, 50  $\mu\text{l}$  of anti-CU antiserum diluted 1:1000 in PBS containing 0.25% (w/v) gelatine and 0.2% (v/v) Tween 20 were added. Rabbit preimmune serum was used in control wells. Plates were incubated overnight at 4°C and washed three times with PBS. Fifty  $\mu\text{l}$ /well of goat anti rabbit IgG-peroxidase conjugate (Sigma-Aldrich, Milan, Italy) diluted 1:2000 in PBS containing 0.25% (w/v) gelatine and 0.2% (v/v) Tween 20 were added and allowed to incubate for 2 h at 37°C. After three washes with PBS, 150  $\mu\text{l}$ /well of substrate

solution (0.4 mg ml<sup>-1</sup> *o*-phenylenediammine dihydrochloride; 0.012% (v/v) H<sub>2</sub>O<sub>2</sub> in 0.1 M citrate-phosphate buffer pH 5.0) were added. The A<sub>492</sub> was measured after 5 min of incubation with a Titertek Multiskan Plus MKII plate reader (Flow Laboratories Inc., McLean, VA, USA). The concentration of CU in individual samples was determined by comparing the mean A<sub>492</sub> obtained for triplicate wells to a standard curve using linear regression analysis. Standard calibration curves had a linear correlation coefficient  $\geq 0.96$  using purified CU over a concentration range of 0.1-30  $\mu\text{g ml}^{-1}$ . The conditions of the assay were optimised to have an A<sub>492</sub> of about 0.5 for the wells containing the highest concentration of CU. Slopes similar to that of standard curves were obtained with serial dilution curves of samples. The sensitivity of the ELISA for CU was of 100 ng ml<sup>-1</sup>. Negative samples always yielded an A<sub>492</sub>  $\leq 0.020$ .


**Statistical analysis.** Data of CU production by *O. ulmi* and *O. novo-ulmi* EAN and NAN were analysed by ANOVA, fixed model. Homogeneous groups were identified by means of Tukey HSD test.

## RESULTS

CU binds to elm cells and determines mortality in a dose-dependent manner. Preliminary experiments were performed in order to find the optimal conditions, in terms of cell concentration and incubation time, to study the binding of CU to elm cells. In these experiments, an amount of CU (5  $\mu\text{M}$ ) sufficient to cause wilt of elm cuttings (Scala *et al.*, 1994) was incubated with suspensions of elm cells for different periods of time. Daily measurements of unbound CU revealed a slow and gradual decrease in CU concentration paralleled by an increase in fluorescence of elm cells in IF assays. On the basis of these results we found that a three-day-incubation was suitable for studying the binding parameters of CU to elm cells and chose CU concentration range and cell density to be used for subsequent experiments.

Viability of cells was measured on a DED susceptible (C06) and a moderately resistant (196/6) elm clone in the presence of increasing concentrations of CU. For both clones a drastic decrease (about 75%) in the percentage of viable cells was observed at CU concentrations above 2  $\mu\text{M}$ . At CU concentrations above 8  $\mu\text{M}$  the mortality of elm cells, as calculated with the Evans blue assay, increased to over 90%. No differences in sensitivity to the cytotoxic effect of CU was found between cells of the two elm clones tested. Binding of CU to cells of both clones, as revealed with fluorescein-labeled anti CU antibodies, progressively increased with CU concentration. Cells appeared faintly fluorescent starting at 0.3  $\mu\text{M}$  CU. At the higher CU concentrations tested, which determined nearly 100% mortality, the fluorescence was intense. An increasing relative fluorescence index was always associated with decreases in cell


viability, indicating a dose-dependent effect of bound CU on elm cell mortality. When the effect of purified CU was tested on fennel or pear cells no fluorescence neither cytotoxicity were observed. These observations give a further indication that CU binding is specific for elm cells and that its cytotoxic effect is mediated by binding (Fig. 1).



**Fig. 1.** Effect of cerato-ulmin on cell viability (open triangles) and relative fluorescence index (solid triangles) of immunostained elm cells of a Dutch elm disease susceptible (C06) and a moderately resistant (196/6) elm clone as a function of increasing cerato-ulmin concentration in 10 ml of a  $5 \cdot 10^4$  ml<sup>-1</sup> cell suspension after a three day incubation. The anti-cerato-ulmin antibodies were visualized by incubating with a fluorescein-conjugated antirabbit IgG antiserum and viewed under a fluorescence microscope. Data on cell survival are the mean  $\pm$  SE of values obtained from samples collected from three replicates obtained in two independent experiments. IF data are based on observation of cells present in 30 microscopy fields in three independent experiments.

The amount of bound CU after 3 days of incubation in the presence of elm cells was determined by semi-quantitative ELISA in cell extracts. Fig. 2 reports the amount of CU bound to elm clones C06 and 196/6 cells, as a function of the initial concentration of CU added to the culture medium. The equation of the saturation binding curve indicated that the maximum quantity of bound CU was  $351.92 \pm 16.56$  ( $R_2 = 0.9924$ ) pmoles mg<sup>-1</sup> d.wt. for the clone C06 and  $355.59 \pm 23.82$  ( $R_2 = 0.9860$ ) for the clone 196/6. The equilibrium dissociation constant was  $7.09 \pm 0.89$   $\mu\text{M}$  CU for the clone C06 and  $8.08 \pm 1.39$   $\mu\text{M}$  CU for the clone 196/6. The Scatchard plot of the saturation binding curve data (not shown), demonstrated that a linear regression was obtained with the same transformed data.

Both fennel and pear cells did not display fluorescence after a 6 day incubation with purified CU and CU concentration in culture fluids did not significantly decrease over the whole period of incubation.



**Fig. 2.** Cerato-ulmin binding to cells of elm clones C06 and 196/6 cells as a function of increasing cerato-ulmin concentration in 10 ml of a  $5 \cdot 10^4$  ml<sup>-1</sup> cell suspension after a three day incubation. Data regarding cerato-ulmin binding are the mean  $\pm$  SE of values obtained from samples collected from three replicates obtained in two independent ELISA determinations.

**Elm cells support growth of *Ophiostoma* fungi and CU production in dual cultures.** Table 1 shows growth and CU production by *O. novo-ulmi* ssp. *americana* isolate H328 and *O. ulmi* isolate E2 in *Ophiostoma*-elm cell dual cultures, prepared utilizing two different liquid media: a medium routinely used for cultivation of *Ophiostoma* (Takai, 1978; abbreviated as TK in Table 1) and a medium routinely used for plant cell cultures

(Murashige & Skoog, abbreviated as MS). Previous work (Scala, 1997) showed that MS only allows limited growth of *Ophiostoma* fungi. Addition of yeast extract (=YE) determines a quite abundant growth of *Ophiostoma* fungi and CU production. In the absence of YE in TK medium (TK<sub>0</sub>) very poor fungal growth occurs and that at least 0.3 g l<sup>-1</sup> of YE are necessary to support both fungal growth and CU production (Scala, 1997).

In TK<sub>0</sub>, the addition of elm cells of either a DED susceptible (C06) and a moderately resistant (196/6) clone to TK<sub>0</sub> supported growth of *Ophiostoma* fungi and CU production, thus fully replacing the effect of YE. In MS the presence of elm cells in MS did not cause a significant effect on fungal growth. The density of blastoconidia reached by both fungi was, in nearly all cases, significantly lower in MS-based than in TK-based media.

CU production was strongly influenced by the culture media. *O. ulmi* did not produce detectable amounts of CU in TK<sub>0</sub>. The addition of C06 or 196/6 elm cells or fennel cells, but not pear cells to TK<sub>0</sub> or MS determined production of CU to levels similar to those observed in TK or in TK<sub>0.3</sub>YE. *O. novo-ulmi* produced no or very low amounts of CU in TK<sub>0</sub> and MS, whereas in TK<sub>0.3</sub>YE and in MS<sub>2</sub>YE *O. novo-ulmi* produced relatively large quantities of CU, not significantly different from those produced in the standard medium TK. The addition of elm cells of both clones or fennel cells, but not pear cells to TK<sub>0</sub> and MS caused marked increases in CU production by *O. novo-ulmi*, even if differences were not statistically significant for MS-based media. IF tests performed on plant cells after 6 days of co-culture (Table 1) indicated that CU produced in dual cultures bound to elm but not to fennel or pear cells, giving a

**Table 1.** Cell density and cerato-ulmin concentration, as assayed by the turbidimetric method and expressed as cerato-ulmin production index, of *O. novo-ulmi* H328 and *O. ulmi* E2 after 6 days of growth in different Takai- or Murashige & Skoog-based media. Relative IF index (on the basis of a 0-3 arbitrary scale, see Materials and Methods) of cells determined on elm, fennel or pear cells are also indicated. Data are the mean  $\pm$  SE of values obtained from samples collected from three replicates obtained in two independent experiments. Values marked with different small letters are significantly different within TK-based media. Values marked with different capital letters are significantly different from values obtained in the TK medium (at  $P \leq 0.05$ , according to the Tukey test).

Medium and cultured cells	<i>Ophiostoma novo-ulmi</i> H328			<i>Ophiostoma ulmi</i> E2		
	Cell Density	Cerato-ulmin production	IF Index	Cell Density	Cerato-ulmin production	IF index
TK	61 $\pm$ 11 aA	354 $\pm$ 61 aA	-	60 $\pm$ 8 aA	9 $\pm$ 2 aA	-
TK <sub>0</sub>	0.1 $\pm$ 0.1 b	0b	-	0.3 $\pm$ 0.1 b	0 b	-
TK <sub>0</sub> + 0.3 g l <sup>-1</sup> YE	90 $\pm$ 9 a	315 $\pm$ 41 a	-	90 $\pm$ 6 a	8 $\pm$ 2 a	-
TK <sub>0</sub> + 196/6 elm cells	100 $\pm$ 9 a	270 $\pm$ 54 a	3	80 $\pm$ 11 a	7 $\pm$ 1 a	1
TK <sub>0</sub> + C06 elm cells	94 $\pm$ 4 a	287 $\pm$ 55 a	3	100 $\pm$ 9 a	8 $\pm$ 2.5 a	1
TK <sub>0</sub> + fennel cells	75 $\pm$ 15 a	189 $\pm$ 18 a	0	80 $\pm$ 12 a	7 $\pm$ 0.5 a	0
TK <sub>0</sub> + pear cells	81 $\pm$ 10 a	0b	0	74 $\pm$ 16 a	0 b	0
MS	2 $\pm$ 0.3 B	2 $\pm$ 1 B	-	2 $\pm$ 0.5 C	0 B	-
MS + 2 g l <sup>-1</sup> YE	45 $\pm$ 5 A	299 $\pm$ 68 A	-	20 $\pm$ 2 B	6 $\pm$ 1 A	-
MS + 196/6 elm cells	10 $\pm$ 1 B	44 $\pm$ 2 B	2	5 $\pm$ 1 C	6 $\pm$ 0.5 A	1
MS + C06 elm cells	10 $\pm$ 1 B	354 B	2	5 $\pm$ 1 C	5 $\pm$ 0.5 A	1
MS + fennel cells	9 $\pm$ 4 B	41 $\pm$ 4 B	0	4 $\pm$ 2 C	5 $\pm$ 1 A	0
MS + pear cells	10 $\pm$ 7 B	0 B	0	5 $\pm$ 1 C	0 B	0

**Table 2.** Comparison of the amounts of cerato-ulmin (as assayed by the turbidimetric method and expressed as cerato-ulmin production index) produced in three different TK-based media (see Materials and Methods) by several isolates of *O. novo ulmi* and *O. ulmi* after 6 days of culture. Data are the mean±SE of values obtained from samples collected from three replicates obtained in two independent experiments. Within each line, different letters indicate that values are significantly different at  $P \leq 0.05$ , according to the Tukey test.

Fungal isolates	Cerato-ulmin production		
	TK	TK <sub>0</sub>	TK <sub>0</sub> + elm cells
<i>O. novo ulmi</i>			
ssp. <i>novo-ulmi</i>			
H 328	351±19 a	0 b	271±85 a
H 327	241±17 a	0 b	185±23 a
ssp. <i>americana</i>			
182	220±19 a	0 b	192±73 a
H 162	221±16 a	0 b	154±44 a
H 351	247±46 a	0 b	156±18 a
<i>O. ulmi</i>			
E 2	8±2 a	0 b	6±1 a
R 21	11±2 a	0 b	7±1 a
Yu 99	6±1 a	0 b	5±1 a
Yu 16	4±1 a	0 b	6±1 a

signal proportional to the amount of CU present in the culture medium.

The ability of elm cells to support both fungal growth and CU production was confirmed on a set of isolates representative of *O. novo-ulmi* ssp. *novo-ulmi* and *americana* and *O. ulmi* (Table 2). The positive effect of elm and fennel but not pear cells to support CU production indicates that elm and fennel but not pear cells apparently have an inducer of CU production.

## DISCUSSION

In this paper we report the occurrence of binding and cytotoxicity of CU to elm cells. CU binding to elm but not to fennel and pear cells, two non-hosts to DED pathogens, strongly supports the assumption that CU is a pathogenicity factor for DED fungi. The binding curve obtained with increasing CU concentrations vs. a standard elm cell suspension shows that cells are saturated with about 360 pmoles CU mg<sup>-1</sup> d.wt. elm cells and the  $K_d$  determined for the binding of CU to elm cells ranges between 7.09 and 8.08 µM for a susceptible and a moderately resistant elm clone. The occurrence of similar values in the binding parameters for the two elm clones tested does not indicate, at least in our system, that CU is a virulence factor in DED. The values, however, could be overestimated since in aqueous solution CU molecules can aggregate to each other (Russo *et al.*, 1982). This can explain, at least in part, why the  $K_d$  determined for CU binding to elm cells is much higher than the  $K_d$  determined for some other plant receptor-ligand combinations (*e.g.* Kooman-Gershmans *et al.*,

1996). More investigation is needed to establish if CU simply adsorbs to elm cell surface or if specific receptor molecules are involved. Work is in progress to study the influence of culture conditions on CU binding in terms of temperature, ionic strength and pH. In the present work we could also observe that the binding was not affected by several washes in 0.15M NaCl and distilled water. The intensity of fluorescence of elm cells and their mortality were correlated to the amount of CU added to the incubation mixture. On the basis of our results and previous observation that leaf wilt correlated to the amount of CU in leaf tissues (Scala *et al.*, 1997b), it can be proposed that CU is produced by the DED fungi in xylematic vessels and is then translocated in leaves where it binds to host cells. The binding would be the preliminary step in causing cellular damage and consequent leaf wilting. Mutual molecular interactions between cell surfaces of host plants and pathogens, and associated signalling mechanisms, are considered to play a fundamental role in pathogenesis (Callow *et al.*, 1988). Presence of receptors at the surface of plant cells for fungal signal molecules was proposed as a logical presumption (Ride, 1992), since cell surface receptors are a common theme in cell-cell recognition events (Chapman *et al.*, 1988). Ride (1992) reviewed various plant-pathogen systems, where some indirect or direct evidence for plant cell receptors have been obtained. Binding of elicitor molecules to plasma membrane receptors as a preliminary step in recognition events leading to hypersensitive response has been described for a peptide elicitor of *Phytophthora megasperma* (Nurnberger *et al.*, 1994) and the race-specific AVR9 elicitor of *Cladosporium fulvum* (Kooman-Gershmans *et al.*, 1996). The

characterization of specific receptors for fusicoccin, a phytotoxin produced by the fungus *Fusicoccum amygdali*, a pathogen of almond, peach and related species, and their role in mediating physiological response(s) of host cells to the ligand, has been of a great interest for both plant physiology and plant pathology (Graniti *et al.*, 1995). Protein receptor for victorin, a host selective toxin produced by the oat pathogen *Cochliobolus victoriae* has also been extensively characterised (Wolpert *et al.*, 1994). In both cases, the study of toxin receptors contributed to a better understanding to the mechanism of action of the toxin and/or the physiological response(s) to the ligand. The occurrence and significance of elm cell receptor(s) for CU in the pathogenic process of DED and in disease symptoms expression, needs more evidence to be elucidated.

On the basis of our results we cannot, at present, exclude that elm cells could lack a specific receptor for CU and that cellular damage could simply derive from the deposition and adhesion of one or more layers of CU on the external surface of elm cell walls, hampering cellular permeability. Experiments with structural analogues of CU should give more information about the type of interaction between CU and elm cells. It also has to be mentioned that CU shares with  $\beta$ -amyloid, a protein causing a number of neurological and degenerative diseases by deposition on cells, the ability to self-assemble and to determine the formation of aggregates (Lorenzo and Yankner, 1994).

Our data confirm Takai's results (1978) that cultural conditions influence CU production. The addition of yeast extract was essential for CU production in both TK<sub>0</sub> e MS. We found that elm cells fully replaced yeast extract in TK<sub>0</sub>, by supporting both abundant fungal growth and CU production. In the presence of elm cells, *O. novo-ulmi* produced quantities of CU always higher than those of *O. ulmi*. This result was obtained by testing several isolates belonging to both subspecies of *O. novo-ulmi* and to *O. ulmi*, reflecting the behaviour of the two species observed in Takai-based media (Brasier, 1991; Scala *et al.*, 1997a) and *in planta* (Scala *et al.*, 1997b). In MS the effect of elm cells on CU production was much less pronounced than in Takai-based media, but still evident. It should be considered that MS is an unusual medium for growing fungi, since it has been formulated for *in vitro* growth of plant callus and cell cultures.

In an attempt to establish a pathogenicity test alternative to whole plant inoculation, Scala *et al.* (1996) found no differences between the degree of colonization of either DED pathogens of calli derived from a susceptible (*U. carpinifolia*) and a resistant (*U. pumila*) elm species. Nonetheless, calli of the susceptible species were more subjected to cell plasmolysis than those of the resistant species. In our experimental system, which also considers elm clones having different levels of susceptibility to DED pathogens, no differences were found in sensitivity to CU. Presumably, during growth

on calli, factors other than CU could be responsible for the differential sensitivity to plasmolysis observed by Scala *et al.* (1996).

Both fennel and pear cells were able to support fungal growth, but only fennel cells induced CU production, indicating that, in the absence of a source of organic nutrients like yeast extract, CU production is not constitutive and does not depend on a elm-specific inducer molecule. More investigation is needed to establish if pear cell culture media lack one or more metabolites necessary for CU production or contain specific suppressors.

#### ACKNOWLEDGEMENTS

The authors gratefully acknowledge Prof. Clive M. Brasier (Forest Research Station, Alice Holt Lodge, Farnham, UK) and Prof. L. Mittempergher (Istituto per la Protezione delle Piante, CNR, Firenze, Italy) for providing the fungal isolates used in this study. Thanks are due to Dr. Silvia Schiff, Dipartimento di Biologia vegetale, Università di Firenze, Italy, for providing actively proliferating *in vitro* fennel and pear callus cultures. This research was supported by the Italian Ministry for Education and Research (Programme PRIN 1999 "Adromicosi e biomolecole").

#### REFERENCES

- Baker C.J., Mock N.M., 1994. An improved method for monitoring cell death in cell suspension and leaf disc assays using Evan's blue. *Plant Cell, Tissue and Organ Culture* **39**: 7-12.
- Bowden C.G., Hintz W.F., Jeng R., Hubbes M., Horgen P.A., 1994. Isolation and characterisation of the cerato-ulmin toxin gene of the Dutch elm disease pathogen, *Ophiostoma ulmi*. *Current Genetics* **25**: 323-329.
- Bowden C.G., Smalley E., Guries R.P., Hubbes M., Temple B., Horgen P.A., 1996. Lack of association between cerato-ulmin production and virulence in *Ophiostoma novo-ulmi*. *Molecular Plant-Microbe Interactions* **9**: 556-564.
- Brasier C.M., 1991. *Ophiostoma novo-ulmi* sp. nov., causative agent of current Dutch elm disease pandemics. *Mycopathologia* **115**: 151-156.
- Brasier C.M., Mehrotra, M.D., 1995. *Ophiostoma himal-ulmi* sp. nov., a new species of Dutch elm disease fungus endemic to the Himalayas. *Mycological Research* **99**: 205-215.
- Brasier C.M., Kirk S., Tegli S., 1995. Naturally occurring non cerato-ulmin producing mutants of *Ophiostoma novo-ulmi* are pathogenic but lack aerial mycelium. *Mycological Research* **99**: 436-440.
- Brasier C.M., Kirk S.A., 2001. Designation of the EAN and NAN races of *Ophiostoma novo-ulmi* as subspecies: their perithecial size differences and geographical distributions. *Mycological Research* **105**: 547-554.

- Brasier C.M., Takai S., Nordin J.H., Richards W.C., 1990. Differences in cerato-ulmin production between the EAN, NAN and non-aggressive subgroups of *Ophiostoma ulmi*. *Plant Pathology* **39**: 231-236.
- Callow J.A., Ray T., Estrada-Garcia T.M., Green J.R., 1988. Molecular signals in plant cell recognition. In: Scannerini S., Smith D., Bonfante-Fasolo P., Gianinazzi-Pearson V. (eds). NATO ASI Series, Series H; Cell Biology, Vol. 17. Cell to cell signals in plant, animal and microbial symbiosis, pp. 167-182. Springer-Verlag, Berlin, Germany.
- Chapman G.P., Ainsworth C.C., Chatham C.J., 1988. Eukaryote cell recognition. Concepts and model systems. Cambridge University Press, Cambridge, UK.
- Del Sorbo G., Scala F., Parrella G., Lorito M., Comparini C., Ruocco M., Scala A., 2000. Functional expression of the gene *cu*, encoding the phytotoxic hydrophobin cerato-ulmin, enables *Ophiostoma quercus*, a non-pathogen on elm, to cause symptoms of Dutch elm disease. *Molecular Plant-Microbe Interactions* **13**: 43-53.
- Del Sorbo G., Scala A., Scala F., Tegli, S., 2002. Cerato-ulmin, a toxin produced by the pathogens of the Dutch elm disease. In: Upadhyay R.K. (ed.). Advances in microbial toxin research and its biotechnological exploitation, pp. 93-104. Kluwer Academic Plenum Publishers, New York, USA.
- Et Touil A., Brasier, C.M., Bernier, L., 1999. Localization of a pathogenicity gene in *Ophiostoma novo-ulmi* and evidence that it may be introgressed from *O. ulmi*. *Molecular Plant-Microbe Interactions* **12**: 6-15.
- Gilchrist D.G., 1983. Molecular modes of action. In: Daly J.M., Deverall B.J. (eds.). Toxins and plant pathogenesis, pp. 81-136. Academic Press, New York, USA.
- Graniti A., Ballio A., Marrè E., 1995. *Fusicoccum (Phomopsis) amygdali*. In: Kohmoto K., Singh U.S., Singh R.P. (eds). Pathogenesis and host specificity in plant diseases: histological, biochemical, genetic and molecular bases. Vol. II, pp. 103-117. Pergamon, Oxford, UK.
- Kooman-Gershmam M., Honee G., Bonnema G., De Wit P.J.G.M., 1996. A high affinity binding site for the AVR9 peptide elicitor of *Cladosporium fulvum* is present on plasma membranes of tomato and other solanaceous plants. *The Plant Cell* **8**: 929-938.
- Lorenzo A., Yanker, B.A., 1994.  $\beta$ -amyloid neurotoxicity requires fibril formation and is inhibited by congo red. *Proceedings of the National Academy of Science USA* **91**: 12243-12247.
- Mitchell R., 1981. Structure: bacterial. In: Durbin R. (ed.) Toxins in Plant Disease, pp. 262-293. Academic Press, New York, USA.
- Nordin J.H., Mason T.L., Smith L.L., Willmann P.A., Richards W.C., Takai S., 1987. Use of Enzyme-Linked Immunosorbent Assay with murine ascitic antibodies to screen micro-organisms for production of cerato-ulmin, a toxin of *Ceratocystis ulmi*. *Phytopathology* **77**: 96-100.
- Nürnberg T., Nennstiel D., Jabs T., Sacks W.R., Hahlbrock K., Scheel D., 1994. High affinity binding of a fungal oligopeptide elicitor to parsley plasma membranes triggers multiple defense responses. *Cell* **78**: 449-460.
- Richards W.C., 1993. Cerato-ulmin: a unique wilt toxin of instrumental significance in the development of Dutch elm disease. In: Sticklen M.B., Sherald J.L. (eds.). Dutch Elm Disease Research, Cellular and Molecular Approaches, pp. 152-170. Springer-Verlag, New York, USA.
- Richards W.C., Takai S., 1973. Novel technique for isolating microstructures present in shake cultures of the fungus *Ceratocystis ulmi*. *Journal of Applied Microbiology* **26**: 443-444.
- Richards W.C., Takai S., 1984. Characterization of the toxicity of cerato-ulmin, the toxin of Dutch elm disease. *Canadian Journal of Plant Pathology* **6**: 291-298.
- Richards W.C., Takai S., 1988. Production of cerato-ulmin in white elm following artificial inoculation with *Ceratocystis ulmi*. *Physiological and Molecular Plant Pathology* **33**: 279-283.
- Ride J.P., 1992. Signals in fungus-plant basic incompatibility. In: Callow J.A., Green J.R. (eds.). Perspectives in plant cell recognition, pp. 213-237. Cambridge University Press, Cambridge, UK.
- Russo P.S., Blum F.D., Ipsen J.D., Abul-Hajj Y.J., Miller W.G., 1982. The surface activity of the phytotoxin cerato-ulmin. *Canadian Journal of Botany* **60**: 1414-1422.
- Scala A. 1997. Elaboration of a nutritive substrate able to support an *in vitro* *Ophiostoma novo-ulmi*- or *O. ulmi*-elm cell suspensions dual culture and cerato-ulmin production. In: *Proceedings of the 10th Congress of the Mediterranean Phytopathological Union, Montpellier, June 1-5, 4*.
- Scala A., Tegli S., Comparini C., Mittempergher L., Scala F., Del Sorbo G., 1994. Influence of fungal inoculum on cerato-ulmin production; purification of cerato-ulmin and detection in elm sucker cuttings. *Petria* **4**: 57-67.
- Scala A., Comparini C., Tegli, S., Ragazzi A., Dellavalle I., Mittempergher L., Lagazio C., 1996. *In vitro* analysis of the interaction between elm and *Ophiostoma ulmi* or *O. novo-ulmi*. *Petria* **6**: 91-103.
- Scala F., Bertelli E., Coppola L., Del Sorbo G., Tegli S., Scala A., 1997a. Comparative determination of cerato-ulmin on cell surface and in mycelial extracts of pathogenic and non-pathogenic *Ophiostoma* species. *Mycological Research* **101**: 829-834.
- Scala A., Pattuelli M., Coppola L., Guastini M., Tegli S., Del Sorbo G., Mittempergher L., Scala F., 1997b. Dutch elm disease progression and quantitative determination of cerato-ulmin in leaves, stems and branches of elms inoculated with *Ophiostoma novo-ulmi* and *O. ulmi*. *Physiological and Molecular Plant Pathology* **50**: 349-360.
- Stringer M.A., Timberlake W.T., 1993. Cerato-ulmin, a toxin involved in Dutch elm disease, is a fungal hydrophobin. *The Plant Cell* **5**: 145-146.
- Svircev A.M., Jeng R.S., Hubbes M., 1988. Detection of cerato-ulmin on aggressive isolates of *Ophiostoma ulmi* by immunochemistry and scanning electron microscopy. *Phytopathology* **78**: 322-327.
- Takai S. 1974. Pathogenicity and cerato-ulmin production in *Ceratocystis ulmi*. *Nature* **252**: 124-126.
- Takai S., 1978. Cerato-ulmin, a wilting toxin of *Ceratocystis*

- ulmi*: cultural factors affecting cerato-ulmin production by the fungus. *Journal of Phytopathology* **91**: 147-158.
- Takai S., Hiratsuka Y., 1980. Accumulation of the material containing the toxin cerato-ulmin on the hyphal surface of *Ceratocystis ulmi*. *Canadian Journal of Botany* **58**: 663-668.
- Takai S., Hiratsuka Y., 1984. Scanning electron microscope observations of internal symptoms of white elm following *Ceratocystis ulmi* infection and cerato-ulmin treatment. *Canadian Journal of Botany* **62**: 1365-1371.
- Takai S., Richards W.C., 1978. Cerato-ulmin, a wilting toxin of *Ceratocystis ulmi*: isolation and some properties of cerato-ulmin from the culture of *C. ulmi*. *Journal of Phytopathology* **91**: 129-147.
- Takai S., Richards W.C., Stevenson K.J., 1983. Evidence for involvement of cerato-ulmin, the *Ceratocystis ulmi* toxin, in the development of Dutch elm disease. *Physiological Plant Pathology* **23**: 275-280.
- Tegli S., Comparini C., Giannetti C., Scala A., 1993. Effect of cerato-ulmin on surviving of *in vitro* elm callus and cells. *Giornale Botanico Italiano* **127**: 190-191.
- Tegli S., Comparini C., Giannetti C., Scala A., 1994. Effect of temperature on growth and cerato-ulmin production of *Ophiostoma novo-ulmi* and *O. ulmi*. *Mycological Research* **98**: 408-412.
- Tegli S., Scala A., 1996. Isolation and characterisation of non cerato-ulmin producing laboratory induced mutants of *Ophiostoma novo-ulmi*. *Mycological Research* **100**: 661-668.
- Temple B., Horgen P.A., Bernier L., Hintz W.E., 1997. Cerato-ulmin, a hydrophobin secreted by the causal agents of Dutch elm disease, is a parasitic fitness factor. *Fungal Genetics and Biology* **22**: 39-53.
- Wolpert T.J., Navarre D.A., Moore D.L., Macko V., 1994. Identification of the 100-kD victorin binding protein from oats. *The Plant Cell* **6**: 1145-1155.
- Yaguchi M., Puszatai M., Roy C., Surewics W.K., Carey P.R., Stevenson K.J., Richards W.C., Takai S., 1993. Amino acid sequence, infrared and raman spectroscopy of Dutch elm disease toxin cerato-ulmin. In: Sticklen M.B., Sherald J.L. (eds.). *Dutch Elm Disease Research, Cellular and Molecular Approaches*, pp. 152-170. Springer-Verlag, New York, USA.
- Yoder O.C., 1980. Toxins in pathogenesis. *Annual Review of Phytopathology* **18**: 103-129.
- Yoder O.C. 1981. Assay. In: Durbin R. (ed.). *Toxins in Plant Disease*, pp. 45-78. Academic Press, New York, USA.

Received 25 February 2003

Accepted 2 July 2003

